

# VIREX



Avian Influenza  
**FREE ZONE**

Poultry  
Farming

**A broad spectrum powder  
disinfectant, effective  
against bacteria, yeasts  
and viruses for surfaces,  
equipment, footbaths and  
animal transport vehicles.**




# VIREX



## ADVANTAGES

- A bactericidal, yeasticidal and virucidal powder disinfectant
- Active chlorine generated from precursors
- For the disinfection of livestock premises, animal breeding equipment and materials, animal transport vehicles and the disinfection of boots in farms.
- Approved for Avian Influenza, Diseases of Poultry & General Orders



## Information & Preparation for Application

Pre-clean surfaces, equipment, materials, animal transport, vehicles, wheels and boots to be disinfected.

Prepare a dilution of VIREX at the recommended rate:

- Bactericidal and Virucidal: 1%
- Yeasticidal: 3%

Minimum Contact Time: 30 minutes. Temperature of use: 10°C minimum

## Spray Application:


Spray the diluted solution on the breeding surfaces, equipment, animal transport vehicles at an average rate of 300 ml/m<sup>2</sup> up to the limit of runoff.

## Dip Application, Boot bath application:

Dip the equipment, materials, wheels and boots in the VIREX solution.



Spray



Dip

### BACTERICIDAL EFFICACY

- **EN 14349: 30 MIN. / 10°C / 0,5% / high soiling** (*Enterococcus hirae*, *Proteus vulgaris*, *Pseudomonas aeruginosa*)
- **EN 14349: 30 MIN. / 10°C / 1% / low soiling** (*Staphylococcus aureus*)
- **EN 1656: 5 MIN. / 10°C / 0,5% / low soiling** (*Enterococcus hirae*, *Proteus vulgaris*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*)

### YEASTICIDAL EFFICACY

- **EN 1657: 5 MIN. / 10°C / 1% / low soiling** (*Candida albicans*)
- **EN 16438: 30 MIN. / 10°C / 3% / low soiling** (*Candida albicans*)

### VIRUCIDAL EFFICACY

- **1%** (*Bovine Enterovirus type 1 (ECBO)*, *African swine fever virus (ASF)*)
- **EN 14675: 30 MIN. / 10°C / low soiling /**
- **EN 14675: 30 MIN. / 10°C / high soiling**
- **0,2%** (*Porcine respiratory & Reproductive Syndrome virus (Lelystad strain)*)
- **0,5%** (*Aujesky's disease virus (strain NIA1)*, *Avian Reovirus (Uchida strain)*, *Bovine parainfluenza type 3 virus (strain 125)*, *Bovine viral diarrhea virus (strain NADL)*, *Chicken Anaemia Virus (strain Cux)*, *Feline Calicivirus (FeCV) strain F9*, *Infectious Bovine Rhinotracheitis virus (JCF strain)*, *Infectious Bursal Disease Virus (strain G13)*, *Infectious Pancreatic Necrosis virus (strain A2, Sp)*, *Porcine Parvovirus (strain 59E)*)
- **1%** (*Avian Influenza (A/Turkey/Eng/N28/73) (H5N1)*, *Egg Drop Syndrome Virus (strain 127)*, *Porcine circovirus type 2 (strain 1010)*)
- **2%** (*Bovine Enterovirus type 1 (strain VG/5/27)*, *Teschen virus (strain F65)*)
- **EN 14675: 4°C**
- **0.5% / 30 MIN. / high soiling:** (*Infectious Pancreatic Necrosis virus (strain A2, Sp)*)
- **1% / 5 MIN. / low soiling:** (*African swine fever virus (ASF)*)
- **1.5% / 30 MIN. / low soiling:** *Avian Influenza Virus*
- **1.5% / 30 MIN. / low soiling:** *Bovine Enterovirus type 1 (ECBO)*

#### DEFRA

**At a dilution rate of 1g disinfectant plus 100ml water**  
*Avian Influenza Order, Influenza of Avian Origin in Mammals, Diseases of Poultry Order, Swine Vesicular Disease*

**At a dilution rate of 1g disinfectant plus 1300ml water**  
*Foot and Mouth Disease*

Active substances: Troclosene sodium + Bis(peroxymonosulfate)bis(sulfate) of pentapotassium  
pH at 10g/l: ≈ 2

Storage: Keep in a clean, cool and well-ventilated place away from sources of heat and intense light.

Shelf Life: 2 years from date of manufacture